



Akebia Initiates Vafseo® (vadadustat) Post-Marketing Study in Conjunction with Large Dialysis Organization

August 4, 2025

VOCAL, an open-label active-controlled study, is being conducted at DaVita dialysis clinics to evaluate benefits of treating patients with Vafseo three times a week

CAMBRIDGE, Mass., Aug. 04, 2025 (GLOBE NEWSWIRE) -- [Akebia Therapeutics®, Inc.](#) (Nasdaq: AKBA), a biopharmaceutical company with the purpose to better the lives of people impacted by kidney disease, today announced the initiation of a post-marketing study to identify additional potential benefits of Vafseo® (vadadustat). The VOCAL trial, conducted within DaVita clinics, aims to generate data to evaluate the efficacy and safety of three times per week (TIW) dosing of vadadustat compared to standard of care erythropoiesis-stimulating agents (ESA) in patients with anemia of CKD receiving in-center hemodialysis.

"Our team and partners within the dialysis community share a commitment to improving patient care," said Steven K. Burke, M.D., Chief Medical Officer of Akebia. "The VOCAL trial, conducted within DaVita clinics, enables us to continue building the body of evidence on the effectiveness of Vafseo when dosed three times a week as well as to understand the potential additional benefits of Vafseo treatment for anemia in CKD patients."

The VOCAL trial is expected to enroll approximately 350 patients across 18 DaVita hemodialysis clinics. This open-label trial will employ 1:1 randomization. Patients will participate in the study for up to 33 weeks including screening, treatment and safety follow-up. The primary endpoint of the study is change in hemoglobin, and secondary endpoints include number of patients reporting treatment-emergent serious adverse events, proportion of patients within target hemoglobin range and proportion of patients receiving RBC transfusions, in each case, compared to ESA treatment.

The VOCAL trial will also include a sub-study of approximately 28 patients from three clinics who will have blood samples collected and analyzed over the course of a 6-month treatment period. One major factor contributing to the anemia of CKD patients is reduced red blood cell (RBC) quality. Working in consultation with Vitalant Research Institute, the intent of the sub-study is to better understand the impact of Vafseo on RBC phenotypes (e.g. deformability, resistance to oxidative stress, metabolomics) compared to ESA treatment.

"DaVita is committed to be at the forefront of innovation by evaluating novel treatments that improve clinical practice and benefit patient care," said Dr. Steven Brunelli, MD, MSCE, Vice President and Medical Director of DaVita Outcomes Research. "The VOCAL trial evaluates Vafseo as a potential new standard of care in a real-world setting and explores additional potential benefits for patients."

Vafseo was approved by the U.S. Food and Drug Administration in March 2024 for the treatment of anemia due to chronic kidney disease (CKD) in adults who have been receiving dialysis for at least three months, and the product became available in the U.S. in January 2025.

About Akebia Therapeutics

Akebia Therapeutics, Inc. is a fully integrated biopharmaceutical company with the purpose to better the lives of people impacted by kidney disease. Akebia was founded in 2007 and is headquartered in Cambridge, Massachusetts. For more information, please visit our website at www.akebia.com, which does not form a part of this release.

About Vafseo® (vadadustat) tablets

Vafseo® (vadadustat) tablets is a once-daily oral hypoxia-inducible factor prolyl hydroxylase inhibitor that activates the physiologic response to hypoxia to stimulate endogenous production of erythropoietin, increasing hemoglobin and red blood cell production to manage anemia. Vafseo is approved for use in 37 countries.

INDICATION

VAFSEO is indicated for the treatment of anemia due to chronic kidney disease (CKD) in adults who have been receiving dialysis for at least three months.

Limitations of Use

- VAFSEO has not been shown to improve quality of life, fatigue, or patient well-being.
- VAFSEO is not indicated for use:
 - As a substitute for red blood cell transfusions in patients who require immediate correction of anemia.
 - In patients with anemia due to CKD not on dialysis.

IMPORTANT SAFETY INFORMATION about VAFSEO (vadadustat) tablets

WARNING: INCREASED RISK OF DEATH, MYOCARDIAL INFARCTION, STROKE, VENOUS THROMBOEMBOLISM, and THROMBOSIS OF

VASCULAR ACCESS.

VPFSEO increases the risk of thrombotic vascular events, including major adverse cardiovascular events (MACE).

Targeting a hemoglobin level greater than 11 g/dL is expected to further increase the risk of death and arterial and venous thrombotic events, as occurs with erythropoietin stimulating agents (ESAs), which also increase erythropoietin levels.

No trial has identified a hemoglobin target level, dose of VPFSEO, or dosing strategy that does not increase these risks.

Use the lowest dose of VPFSEO sufficient to reduce the need for red blood cell transfusions.

CONTRAINDICATIONS

- Known hypersensitivity to VPFSEO or any of its components
- Uncontrolled hypertension

WARNINGS AND PRECAUTIONS

• Increased Risk of Death, Myocardial Infarction (MI), Stroke, Venous Thromboembolism, and Thrombosis of Vascular Access

A rise in hemoglobin (Hb) levels greater than 1 g/dL over 2 weeks can increase these risks. Avoid in patients with a history of MI, cerebrovascular event, or acute coronary syndrome within the 3 months prior to starting VPFSEO. Targeting a Hb level of greater than 11 g/dL is expected to further increase the risk of death and arterial and venous thrombotic events. Use the lowest effective dose to reduce the need for red blood cell (RBC) transfusions. Adhere to dosing and Hb monitoring recommendations to avoid excessive erythropoiesis.

• Hepatotoxicity

Hepatocellular injury attributed to VPFSEO was reported in less than 1% of patients, including one severe case with jaundice. Elevated serum ALT, AST, and bilirubin levels were observed in 1.8%, 1.8%, and 0.3% of CKD patients treated with VPFSEO, respectively. Measure ALT, AST, and bilirubin before treatment and monthly for the first 6 months, then as clinically indicated. Discontinue VPFSEO if ALT or AST is persistently elevated or accompanied by elevated bilirubin. Not recommended in patients with cirrhosis or active, acute liver disease.

• Hypertension

Worsening of hypertension was reported in 14% of VPFSEO and 17% of darbepoetin alfa patients. Serious worsening of hypertension was reported in 2.7% of VPFSEO and 3% of darbepoetin alfa patients. Cases of hypertensive crisis, including hypertensive encephalopathy and seizures, have also been reported in patients receiving VPFSEO. Monitor blood pressure. Adjust anti-hypertensive therapy as needed.

• Seizures

Seizures occurred in 1.6% of VPFSEO and 1.6% of darbepoetin alfa patients. Monitor for new-onset seizures, premonitory symptoms, or change in seizure frequency.

• Gastrointestinal (GI) Erosion

Gastric or esophageal erosions occurred in 6.4% of VPFSEO and 5.3% of darbepoetin alfa patients. Serious GI erosions, including GI bleeding and the need for RBC transfusions, were reported in 3.4% of VPFSEO and 3.3% of darbepoetin alfa patients. Consider this risk in patients at increased risk of GI erosion. Advise patients about signs of erosions and GI bleeding and urge them to seek prompt medical care if present.

• Serious Adverse Reactions in Patients with Anemia Due to CKD and Not on Dialysis

The safety of VPFSEO has not been established for the treatment of anemia due to CKD in adults not on dialysis and its use is not recommended in this setting. In large clinical trials in adults with anemia of CKD who were not on dialysis, an increased risk of mortality, stroke, MI, serious acute kidney injury, serious hepatic injury, and serious GI erosions was observed in patients treated with VPFSEO compared to darbepoetin alfa.

• Malignancy

VPFSEO has not been studied and is not recommended in patients with active malignancies. Malignancies were observed in 2.2% of VPFSEO and 3.0% of darbepoetin alfa patients. No evidence of increased carcinogenicity was observed in animal studies.

ADVERSE REACTIONS

- The most common adverse reactions (occurring at $\geq 10\%$) were hypertension and diarrhea.

DRUG INTERACTIONS

- **Iron supplements and iron-containing phosphate binders:** Administer VPFSEO at least 1 hour before products containing iron.

- **Non-iron-containing phosphate binders:** Administer VAFSEO at least 1 hour before or 2 hours after non-iron-containing phosphate binders.
- **BCRP substrates:** Monitor for signs of substrate adverse reactions and consider dose reduction.
- **Statins:** Monitor for statin-related adverse reactions. Limit the daily dose of simvastatin to 20 mg and rosuvastatin to 5 mg.

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** May cause fetal harm.
- **Lactation:** Breastfeeding not recommended until two days after the final dose.
- **Hepatic Impairment:** Not recommended in patients with cirrhosis or active, acute liver disease.

Please note that this information is not comprehensive. Please click [here](#) for the Full Prescribing Information, including **BOXED WARNING** and Medication Guide.

Forward-Looking Statements

Statements in this press release regarding Akebia Therapeutics, Inc.'s ("Akebia's") strategy, plans, prospects, expectations, beliefs, intentions and goals are forward-looking statements within the meaning of the U.S. Private Securities Litigation Reform Act of 1995, as amended, and include, but are not limited to, statements regarding: Akebia's expectations regarding the VOCAL trial, including expectations as to the potential benefits of Vafseo for the treatment of anemia in CKD patients, potential benefits of Vafseo when dosed three times a week and the expected patient enrollment; and Akebia's beliefs and plans to establish Vafseo as the new standard of care for the treatment of anemia due to CKD. The terms "intend," "believe," "plan," "goal," "potential," "anticipate," "estimate," "expect," "future," "will," "continue," derivatives of these words, and similar references are intended to identify forward-looking statements, although not all forward-looking statements contain these identifying words. Actual results, performance or experience may differ materially from those expressed or implied by any forward-looking statement as a result of various risks, uncertainties and other factors, including, but not limited to, risks associated with: the potential therapeutic benefits, safety profile, and effectiveness of Vafseo; the results of preclinical and clinical research; decisions made by health authorities, such as the FDA, with respect to regulatory filings and other interactions; the commercial availability of Vafseo; the potential demand and market potential and acceptance of, as well as coverage and reimbursement related to, Vafseo, including estimates regarding the potential market opportunity; the competitive landscape for Vafseo, including generic entrants and the timing thereof; the ability of Akebia to attract and retain qualified personnel; Akebia's ability to achieve and maintain profitability and to maintain operating expenses consistent with its operating plan; manufacturing, supply chain and quality matters and any recalls, write-downs, impairments or other related consequences or potential consequences; early termination of any of Akebia's collaborations; and changes in the geopolitical environment and uncertainty surrounding U.S. trade policy on tariffs. Other risks and uncertainties include those identified under the heading "Risk Factors" in Akebia's Quarterly Report on Form 10-Q for the quarter ended March 31, 2025, and other filings that Akebia may make with the U.S. Securities and Exchange Commission in the future. These forward-looking statements (except as otherwise noted) speak only as of the date of this press release, and, except as required by law, Akebia does not undertake, and specifically disclaims, any obligation to update any forward-looking statements contained in this press release.

Akebia Therapeutics® and Vafseo® are registered trademarks of Akebia Therapeutics, Inc.

Akebia Therapeutics Contact

Mercedes Carrasco
mcarrasco@akebia.com



Source: Akebia Therapeutics, Inc.